

Online-Only Abstract

This article is available online at wileyonlinelibrary.com

Transmission of *Staphylococcus aureus* between mothers and infants in an African setting

F. Schaumburg^{1,2}, A. S. Alabi^{2,3}, G. Mombo-Ngoma^{3,4,5}, H. Kaba^{2,3}, R. M. Zoleko^{2,3,5}, D. A. Diop^{2,3}, J.-R. Mackanga^{2,3}, A. Basra^{2,3,5}, R. Gonzalez⁶, C. Menendez⁶, M. P. Grobusch^{2,3,7}, P. G. Kremsner^{2,3}, R. Köck⁸, G. Peters¹, M. Ramharter^{2,3,9} and K. Becker¹

1) Institute of Medical Microbiology, University Hospital Münster, Münster, Germany, 2) Centre de Recherches Médicales de Lambaréné, Albert Schweitzer Hospital, Lambaréné, Gabon, 3) Institut für Tropenmedizin, Eberhard Karls Universität, Tübingen, Germany, 4) Département de Parasitologie-Mycologie, Université des Sciences de la Santé, Libreville, Gabon, 5) Ngounie Medical Research Centre, Fougamou, Gabon, 6) Barcelona Centre for International Health Research (CRESIB), Hospital Clinic University of Barcelona, Barcelona, Spain, 7) Department of Infectious Diseases, Division of Internal Medicine, Centre of Tropical Medicine and Travel Medicine, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands, 8) Institute of Hygiene, University Hospital Münster, Münster, Germany and 9) Department of Medicine I, Division of Infectious Diseases and Tropical Medicine, Medical University of Vienna, Vienna, Austria

Original Submission: 15 July 2013; **Revised Submission:** 28 August 2013; **Accepted:** 26 September 2013

Editor: G. Lina

Article published online: 18 November 2013

Clin Microbiol Infect 2014; **20**: O390–O396

10.1111/1469-0691.12417

Abstract

Staphylococcus aureus colonization is a risk factor for invasive disease. There is a need to understand *S. aureus* colonization in infancy as the burden of *S. aureus* infections in infants is high. We aimed to investigate the transmission of *S. aureus* between mothers and their newborns during the first year after delivery in an African setting. In a longitudinal cohort study, colonization of Gabonese mother–infant pairs was assessed at delivery and after 1, 9 and 12 months. Swabs were taken from mothers (nares, mammillae) and infants (nares and throat). Isolates were characterized and risk factors for colonization were assessed using a standardized questionnaire. We recruited 311 mothers and 318 infants including seven sets of twins. Maternal and infant colonization rates declined synchronously following a peak after 1 month at 40% (mothers) and 42% (infants). Maternal colonization was a risk factor for *S. aureus* carriage in infants. Based on *spa* typing, direct mother-to-infant transmission was evident in 5.6%. Of all methicillin-resistant isolates ($n = 9$), 44.4% were related to the USA300 clone; 56.7% ($n = 261$) of all *S. aureus* carried Panton–Valentine leukocidin encoding genes. Direct mother-to-infant transmission was rare and cannot explain the increase of carriage in infants within the first month. A transmission from external sources is likely and challenges the *S. aureus* infection control in newborns and infants in an African setting. The detection of USA300-related MRSA fuels the concern about the spread of this clone in Central Africa.